

Environmental Information Sheet

Boogie[®] xpro[®] MAPP 15061



An emulsifiable concentrate formulation containing 50 g/L (5.1% w/w) bixafen and 100 g/L (10.1% w/w) prothioconazole and 250 g/L (25.3% w/w) spiroxamine.

BOOGIE xpro is a mixture of a carboxamide, triazolinthione and spiroketoalamine fungicides recommended for control of a wide range of diseases on winter and spring wheat, triticale and winter rye.

Maximum individual dose: Winter and spring wheat, triticale and winter rye: 1.5 litres product per hectare

Maximum number of treatments: Two per crop

Latest time of application: Before grain milky ripe stage

Section	Profile
<p>1. WILDLIFE</p> <p>Mammals and Birds</p>	<p>No risk management is necessary to protect mammals and birds.</p> <p>Boogie xpro is not classified as <i>“Harmful to game, wild birds and animals.”</i></p> <p>Boogie xpro is of low toxicity to mammals and birds. The risk to wild mammals and birds grazing on treated areas is low, as is the risk due to exposure from other routes, e.g. consumption of earthworms or other invertebrates such as insects.</p>
<p>2. BEES</p>	<p>No risk management necessary.</p> <p>Boogie xpro is of low risk to honeybees</p>
<p>3. NON TARGET INSECTS AND OTHER ARTHROPODS</p>	<p>No risk management necessary. Boogie xpro poses a low risk to a range of arthropod species commonly found in and around treated fields, e.g. ground beetles, lady birds and aphid parasitoids.</p>
<p>4. AQUATIC LIFE</p>	<p>Boogie xpro is classified as <i>“Very toxic to aquatic life with long lasting effects.” Do not contaminate water with the product or its container”</i>. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).</p> <p>Risk management is necessary.</p> <p>Boogie xpro can be used safely providing care is taken to prevent spray drift reaching surface waters. The following risk management practices must be carried out in order to ensure that there is adequate protection for aquatic species. DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water</p>

Environmental Information Sheet

Boogie[®] Xpro[®] MAPP 15061



4. AQUATIC LIFE	<i>LERAP Category B. Buffer zone may be reduced</i>
5. SOIL and GROUNDWATER Earthworms Soil Micro-organisms	<p>Bixafen is very persistent in soil and is of low mobility in soil. Prothioconazole is non-persistent in soil and of low mobility in soil. Spiroxamine is moderately persistent in soil and is of low mobility in soil.</p> <p>Therefore, use of Boogie Xpro according to the label presents a low risk to groundwater.</p> <p>Bixafen and prothioconazole and spiroxamine are of low toxicity to earthworms. No risk management is necessary, when Boogie Xpro is used according to the label</p> <p>Boogie Xpro is of low risk to soil micro-organisms. No risk management necessary in order to prevent risks to the processes of soil respiration and nitrogen turnover.</p>
6. NON-TARGET PLANTS	No effects on non target plant species are expected when BoogieXpro is used as recommended.

Use plant protection products safely. Always read the label and product information before use. For further product information including warning phrases and symbols refer to www.cropscience.bayer.co.uk

For further information about the environmental profile of this product contact:

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge
CB4 0WB

Bayer CropScience Assist Technical Help-Line: 0808 1969522

Boogie[®] and Xpro[®] are registered trademarks of Bayer
© Bayer CropScience Limited 2011