

Environmental Information Sheet

FOLICUR® MAPP 16731



An emulsion (oil in water) formulation containing 250 g/l (25.9% w/w) tebuconazole
A broad spectrum systemic fungicide for winter and spring crops of wheat (excluding durum), barley, oats, rye and oilseed rape; also field beans and linseed.

Wheat, barley, rye (winter) and oats: Maximum individual dose: 1.0 L product per hectare
 Maximum total dose: 2.0 L product per hectare (see Other specific restrictions).

Oilseed rape: (a) First application between growth stages BBCH 14 and at or before BBCH 19 Maximum individual dose 0.5 L product per hectare Maximum total dose 0.5 L product per hectare (see Other specific restrictions). **Or**

b) First application at or after BBCH 20. Maximum individual dose: 1.0 L product per hectare
 Maximum total dose: 1.0 L product per hectare (see Other specific restrictions).

Field beans and linseed: Maximum individual dose: 1.0 L product per hectare, Maximum total dose: 1.0 L product per hectare.

Section	Profile
1. WILDLIFE Mammals and Birds	Folicur is not classified as <i>"Harmful to game, wild birds and animals."</i> No risk management is necessary to protect mammals and birds. Folicur is of low/moderate toxicity to mammals and birds. The risk to wild mammals and birds grazing on treated areas is low, as is the risk due to exposure from other routes, e.g. consumption of insects.
2. BEES	No risk management necessary. Folicur is of low risk to honeybees
3. NON TARGET INSECTS AND OTHER ARTHROPODS	No risk management necessary. Folicur poses a low risk to a range of arthropod species commonly found in and around treated fields, e.g. ground beetles, lady birds and aphid parasitoids.
4. AQUATIC LIFE	Folicur is classified as Very toxic to aquatic life with long lasting effects. Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads). Folicur can be used safely providing care is taken to prevent spray drift reaching surface waters. The following risk management practices must be carried out in order to ensure that there is adequate protection for aquatic species. DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water LERAP Category B. Buffer zone may be reduced

Environmental Information Sheet

FOLICUR® MAPP 16731



Section	Profile
5. SOIL and GROUNDWATER	Tebuconazole is moderately persistent in soil, but has low mobility in soil. Use of Folicur according to the label presents a low risk to groundwater.
Earthworms	Tebuconazole is of low toxicity to earthworms. No risk management is necessary, when Folicur is used according to the label.
Soil Micro-organisms	Folicur is of low risk to soil micro-organisms. No risk management necessary in order to prevent risks to the processes of soil respiration and nitrogen turnover.
6. NON-TARGET PLANTS	When used as recommended Folicur is not expected to have adverse effects on non-target plants.

Use plant protection products safely. Always read the label and product information before use. For further product information including warning phrases and symbols refer to www.cropscience.bayer.co.uk

For further information about the environmental profile of this product contact:

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge
CB4 0WB

Bayer CropScience Assist Technical Help-Line: 0808 1969522

Folicur® is a registered trademark of Bayer
© Bayer CropScience Limited 2011