



Please note that physical compatibilities may not be approved tank-mixes. The tests do not check for any adverse crop phytotoxicity or for the biological efficacy of the individual components when applied in a tank-mix. Bayer CropScience gives no warranty and accepts no liability in respect of physical compatibilities; therefore, use is at grower's own risk.

Folicur may be applied in specified tank-mixtures. The matrix charts the recommended Herbicide, PGR and Fungicide compatibilities.

Please note that only tank mixtures selected from a single column and a single row are recommended, e.g. Folicur + Swift SC + chlorothalonil + Eagle.

Fungicides	Fungicides						Herbicides														Plant Growth regulators											
	chlorothalonil SC*	Corbel	Fortress	Tern	Torch Extra	Swift sc	Ally Max SX Max SX	Attribut	Boxer	Capture	Cheetah Super/Triumph + Toil	Chekker	mecoprop-p	Eagle	Grasp + Output	Harmony M SX	Hussar	Oxytril CM	Quantum SX	Sekator	Starane 2	Topik + Actipron	Cerone	Chlormequat	5C Cycocel	Chlormequat + Moddus†	Moddus	Meteor	Meteor + Moddus	Terpal + Agral	Upgrade	
Folicur	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Folicur + Amistar	✓	✓	✓	✓	✓		✓				✓		✓	✓	✓			✓			✓		✓	✓	✓						✓	
Folicur + Comet/Tucana/Vivid	✓	✓	✓	✓	✓								✓	✓	✓			✓			✓										✓	
Folicur + Corbel	✓					✓	✓				✓		✓			✓								✓	✓							
Folicur + Tern	✓					✓																		✓	✓							✓
Folicur + Torch Extra	✓		✓				✓						✓							✓			✓							✓		
Folicur + Swift SC	✓	✓	✓	✓	✓		✓		✓				✓	✓	✓				✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Folicur + Swift SC + chlorothalonil SC*		✓	✓	✓	✓		✓		✓				✓	✓	✓				✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

✓ Compatible	Blank: Not known	* Approved chlorothalonil SC formulations only
* Not compatible	† At rates specified below	



- In addition to the tank-mix recommendations on page 1, Folicur can also be tank-mixed with one additional product from either of the lists of insecticides or micro-nutrients.

Insecticides	Micro-nutrients			
Aphox	Bittersaltz	Headland Jett	Headland Thio-S	YaraVita Mantrac DF
Cypermethrin	Cutonic Copper Ultra	Headland Magnesium	Librel liquid Copper	YaraVita Mantrac 500
Decis	Cutonic Boron 150	Headland Mansion	Librel Liquid Magnesium	YaraVita Sulpur F3000
Fury 10 EW	Cutonic Manganese Ultra	Headland Mn 15%	Librel Liquid Manganese*	Vytel Liquid Boron†
Hallmark with Zeon Technology	Epsom salt†	Headland Root 66	Omex Protein Plus	Vytel Liquid Copper
Mavrik	Headland Boron 15	Headland Spirit	Omex Protein Plus S	Vytel Liquid Magnesium
Sumi-Alpha	Headland Hi-Phos	Headland Stag	YaraVita Coptrel 500	Vytel Liquid Manganese

† Physically compatible with constant agitation. Do not allow to stand in tank for long periods without agitation.

* Add Folicur last

Additional tank mixes				
Folicur + Ally Max SX + chlormequat	Folicur + Corbel + Harmony M	Folicur + Ensign	Folicur + Stronghold	Folicur + Thiovit Jet
Folicur + Ally Max SX + mecoprop-p†	Folicur + Comet + Stomp 2 [†]	Folicur + Fandango	Folicur + Swift SC + Opus	Folicur + Tern + Sanction
Folicur + Ally Max SX + Starane 2	Folicur + Corbel + Ally Max SX + Cheetah Super	Folicur + Flexity	Folicur + Swipe	Folicur + Unix*
Folicur + Amistar + Starane 2	Folicur + Corbel + Ally Max SX + chlormequat	Folicur + Opus	Folicur + Talius	
Folicur + Centaur	Folicur + Corbel + Ally Max SX + mecoprop-p	Folicur + Proline ²⁷⁵	Folicur + Tern + chlormequat + Sanction	
Folicur + chlormequat + Starane 2	Folicur + Cyflamid	Folicur + Punch C	Folicur + Tern + propiconazole	
Folicur + Corbel + chlormequat	Folicur + Dow Shield			

Tank-mix precautions

PGR tank mixtures

Folicur + strobilurin may be applied, before crop GS32, in mixture with one of the following (dose/ha): -
 Moddus 0.2L + chlormequat 1.25L (winter wheat)
 Moddus 0.1L + chlormequat 1.25L (winter barley – light soils or malting crops)
 Moddus 0.2L + chlormequat 1.25L (winter barley – Other soils and feed crops)
 Before using any of the mixtures shown above, first check the manufacturer's latest recommendations for Moddus. DO NOT use Folicur + strobilurin + Moddus + chlormequat on Spring Barley.

Adjuvants

In soft crops avoid the use of tank-mixtures with HBNs or Atlantis WG + biopower. In particular, tank mixtures with EC fungicide formulations as this may result in transitory adverse crop effects.

Sulphonyl Ureas (SU)

All sulphonyl urea herbicides have the potential to cause damage to sensitive crops if residues are left in the spray tank. For this reason sprayer hygiene following the use of SU + chlorothalonil + Folicur is very important. Sprayers should always be thoroughly cleaned out with a proprietary sprayer cleaner immediately after use. This should always take place, even if further cereal crops have been treated by the sprayer between the last application SU mixtures and a subsequent broad leaved crop.

Cheetah Super, Grasp or Topik

Adjuvant oil recommended by the herbicide manufacturer may be added in mixes with Folicur + strobilurin. Do NOT use any of these herbicides in mixes with Folicur + strobilurin + Moddus + chlormequat.

Trace Elements

These products are physically compatible only, there is no information on crop safety available

General Precautions

For any tank-mix it is important that the appropriate manufacturer's recommendations should be consulted and followed, and care taken to ensure that applications are made within the recommendations for each product. First disperse Folicur in the spray tank (follow label 'Mixing' instructions), then add the other product(s), unless specified otherwise.

All mixes should be sprayed immediately and agitated throughout until the spray tank is empty. The user should ensure that all products to be tank-mixed with Folicur are approved for use on the crops which are to be sprayed. Always check individual product labels for any possible restrictions before making a tank-mix recommendation.

Contact Bayer CropScience Limited for full details.

Recent Changes	
13/08/2014	Format update