

Seed
treated
with

ra2ST

Modesto®



Containing 400 g/L (32.5% w/w) clothianidin
and 80 g/L (6.51% w/w) beta-cyfluthrin

Agrichem® HY-PRO Duet

contains 180.47 g/L prochloraz
and 344.98 g/L thiram



Bayer CropScience Limited, 230 Cambridge Science Park
Milton Road, Cambridge CB4 0WB Telephone: 01223 226500

Agrichem (International) Ltd., Industrial Estate, Station Road,
Whittlesey, Peterborough, Cambs PE7 2EY Tel 01733 204019

Seed treatment for winter oilseed rape to improve crop establishment by reducing seedling and foliar damage caused by adult flea beetles, turnip sawfly and aphids and reducing the spread of turnip yellows virus (TuYV formerly BWYV).

The active substance clothianidin is a neo-nicotinoid insecticide and beta-cyfluthrin is a pyrethroid insecticide. Consider resistance management when using subsequent foliar applications. Consult the IRAG website for further information.

PRECAUTIONS

SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND
SUITABLE PROTECTIVE GLOVES when handling contaminated
surfaces and treated seed.

DO NOT HANDLE seed unnecessarily.

DO NOT USE TREATED SEED as food or feed.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE BEEN
USED FOR TREATED SEED for food or feed.

BURY OR REMOVE SPILLAGES.

KEEP TREATED SEED SECURE from people, domestic stock/
pets and wildlife at all times during storage and use.

DANGEROUS TO BEES. To protect bees and pollinating insects
do not drill treated seed where bees are actively foraging.

Do not drill treated seed when flowering weeds are present.

HARMFUL TO GAME AND OTHER WILDLIFE. To protect birds
and wild mammals treated seed should not be left on the soil

surface. Bury or remove spillages.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE
BEEN USED FOR TREATED SEED for food or feed.

WASH HANDS AND EXPOSED SKIN before meals and after
work.

TREATED SEED SHOULD NOT BE BROADCAST.

DO NOT APPLY TREATED SEED FROM THE AIR

DIRECTIONS FOR USE

**Adequate seed drilling equipment shall be used to ensure
a high degree of incorporation in soil, minimisation of
spillage and minimisation of dust emission.**

Treated seed should be stored in a cool, dry, draught
and frost-free place until drilled. It is recommended that
MODESTO treated seed should be sown within 18 months of
seed treatment application.

Treatment with Agrichem HY-PRO Duet may affect the flow
rate of seed through the seed drill. It is important to calibrate
the seed drill before each drilling operation and seed lot.

Follow-up Spray Treatments

If follow-up insecticide sprays are required, the first application
should be with a product of a different chemical group from
the neo-nicotinoids and subsequently alternated.

® Modesto is a Registered Trademark of Bayer

® Agrichem HY-PRO Duet is a Registered Trademark of Agrichem

24 hr Emergency No: 0800 220876