

Environmental Information Sheet

SWIFT® SC MAPP 11227



A 500g/l SC formulation containing trifloxystrobin. A strobilurin fungicide to control a wide range of diseases on winter wheat and winter and spring barley

Maximum application rate: 0.5 litre product/ha.

Maximum no. of applications: 2 per crop per year.

Latest time of application: 35 days before crop harvest

Section	Profile
1. WILDLIFE Mammals and Birds	Swift SC is not classified as <i>"Harmful to game, wild birds and animals."</i> No risk management is necessary to protect mammals and birds. Swift SC is of low toxicity to mammals and birds. The risk to wild mammals and birds grazing on treated areas is low, as is the risk due to exposure from other routes, e.g. consumption of earthworms or other invertebrates such as insects.
2. BEES	No risk management necessary. Swift SC is of low risk to honeybees and there is no requirement to avoid application of the product when bees may be foraging on flowering weeds.
3. NON TARGET INSECTS AND OTHER ARTHROPODS	No risk management necessary. The use of Swift SC does not pose an unacceptable risk to non-target arthropods.
4. AQUATIC LIFE	Swift SC is classified <i>"Very toxic to aquatic life with long lasting effects." Do not contaminate surface waters or ditches with chemical or used container"</i> Swift SC is harmful to certain aquatic organisms and care should be taken to minimise drift when applying Swift SC next to shallow ponds, ditches or slow flowing streams. No risk management necessary. When used according to the label instructions there will not be sufficient contamination of water to present a risk to aquatic life. Not categorised under the LERAP scheme

The
Voluntary
Initiative

February 2017