

Fungicide

INFINITO®

A protectant/systemic fungicide used to control late blight in potatoes.

**A suspension concentrate containing
62.5 g/L (5.53 % w/w) fluopicolide +
625 g/L (55.3 %w/w) propamocarb
hydrochloride**

The (COSHH) Control of Substances
Hazardous to Health Regulations may apply
to the use of this product at work.

Bayer CropScience Ltd
230 Cambridge Science Park
Milton Road, Cambridge, CB4 0WB
Telephone: 01223 226500

**For 24 hour emergency information
contact Bayer CropScience Ltd
Telephone: 00800 1020 3333**

10 L e



GROUP 28 | 43 FUNGICIDES

MAPP 16335 F



GB85419501f rA3b

Safety Information

INFINITO

UFI: 5UQ0-T0CN-300N-SQ2J
 Contains 62.5 g/L (5.53% w/w) fluopicolide and 625 g/L (55.3% w/w) propamocarb hydrochloride.

**Warning**

May cause an allergic skin reaction.

Suspected of damaging the unborn child.
 Very toxic to aquatic life with long lasting effects.
 Wear protective gloves/protective clothing/eye protection/face protection.

If skin irritation or rash occurs: Get medical advice/attention.

IF exposed or concerned: Call a POISON CENTER/doctor/physician.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop:	Potato
Maximum individual dose:	1.6 L product per hectare
Maximum total dose:	6.4 L product per hectare per crop
Latest time of application:	7 days before harvest

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

PROTECT FROM FROST
 SHAKE WELL BEFORE USE

Bayer CropScience Ltd
 230 Cambridge Science Park
 Milton Road, Cambridge, CB4 0WB
 Telephone: 01223 226500

Bayer

**For 24 hour emergency information
 contact Bayer CropScience Ltd
 Telephone: 00800 1020 3333**



To access the **Safety Data Sheet** for this product scan the code or use the link below:

<https://cropscience.bayer.co.uk/our-products/fungicides/infito>

or alternatively contact your supplier

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WASH CONCENTRATE from skin or eyes immediately.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

Environmental Protection

DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect ground water do not apply this or any other product containing fluopicolide at a rate of more than 400 g/ha in any three year period.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This leaflet is approved as part of the label. All instructions on this leaflet and on the label should be read carefully in order to obtain successful results from the use of this product.

RESTRICTIONS

Apply Infinito only to dry foliage. Do not apply Infinito if rainfall or irrigation is imminent. Rainfastness: 1 hour, providing the spray has dried on the leaf.

Avoid drift to areas outside those being sprayed having due regard to the prevailing weather conditions and the spray quality being used.

CROP SPECIFIC INFORMATION

Infinito may be used on all varieties of potato and crops grown for seed.

DISEASES CONTROLLED

When used as part of a full blight protection programme of correctly timed sprays, Infinito gives control of late blight of potatoes (*Phytophthora infestans*).

Infinito will protect tubers from the risk of late blight infection after harvest and reduce tuber blight incidence when used as part of a full blight protection programme.

Resistance management

To reduce the incidence of resistance developing, Infinito should be used in single or block applications in alternation with fungicides from a different cross-resistance group.

TIMING AND DOSE

Applications must be made preventively as part of a blight control programme with regard to blight infection periods or local blight warnings. In the absence of any warnings, applications should commence when the crop meets along the row.

Apply Infinito at 1.2 to 1.6 L/ha on a 7-10 day spray interval, depending on disease pressure. As disease pressure and the risk of late blight infection increases, the dose should be increased and the interval shortened. Irrigated crops should be treated as high risk.

Infinito may be applied at a shorter interval than indicated above should there be an agronomic need as long as all other conditions of use are followed – particularly the maximum number of treatments, maximum total dose and the latest timing.

Infinito should be applied as a protectant treatment. Do not apply as a curative treatment when blight is present in the crop. When blight has become readily visible (1% leaf area destroyed), fungicides have little effect on subsequent development. Stop using Infinito under these conditions.

Tuber blight

Used as part of a full blight protection programme, Infinito will protect tubers from the risk of late blight infection after harvest and reduce tuber blight incidence when applied at the recommended dose and spray intervals from full canopy development to haulm desiccation.

APPLICATION

Infito must be applied to dry foliage using appropriate spray equipment.

Water volume

Apply in 200-400 litres of water per hectare

SPRAY QUALITY

Apply as a MEDIUM quality spray (for details see Boom Sprayers Handbook published by BCPC).

SPRAY OPERATORS INSTRUCTIONS

SHAKE WELL BEFORE USE

1. Wash out sprayer, spray bars and nozzles to ensure that no trace remains of previous chemicals.
2. Check that the nozzle tips are clean, undamaged, of the correct type to produce the recommended spray quality and all of the same size.
3. Start filling the sprayer tank with clean water; check that the agitation is operating correctly. Check dose and volume of application.
4. Calculate the area to be sprayed with the tank load and hence the amount of chemical required. Add the required quantity of product to the water in the sprayer whilst the agitator is operating. Mix only sufficient chemical for immediate use.

5. Add water to the required level and continue agitation until spraying is completed.

6. Ensure that the spray bars are set at the correct height to give a good spray cover, and that alternate spray cones or fans meet just above the crop. Even cover with a well-mixed spray is essential for good results.

7. Commence spraying, being careful to maintain the correct pressure and tractor speed.

8. Wash equipment thoroughly with water and liquid detergent immediately after use. Finally wash out twice with water and drain.

CONTAINER DISPOSAL

Empty containers should be disposed of in a safe way.

DISEASE RESISTANCE

Certain diseases may develop resistance to Bayer CropScience Limited's products. Since such circumstances are beyond our control, Bayer CropScience can accept no liability for any loss or damage whatsoever.

© Infito is a Registered Trade Mark of Bayer

© Bayer CropScience Limited 2024